

BACILLUS CEREUS SCHEMA

for whole genome sequence typing

We are delighted to announce a schema for true whole-genome multi-locus sequence typing (wgMLST) of Bacillus cereus in BioNumerics. The schema brings easy and highly discriminatory detection of subtype- or outbreak-specific markers from whole genome sequencing data to your fingertips.

What is the schema exactly?

Using a selection of 372 annotated, publically available reference genomes capturing the known diversity within *B. cereus*, a pangenomic schema with high discriminatory power was developed. Starting from the reference genomes, our in-house developed schema creation procedure uses a sampling-based multi-reciprocal BLAST procedure to determine those sets of alleles

that make up the stable loci in the pan-genome. A per-locus allele assessment procedure then determines the central prototype allele, and thus the definition of the locus. These loci were further validated using another selection of 589 annotated, publically available genomes. In this way, a total of 30,356 validated loci were added to the schema. Additionally, the 7 classic MLST loci were added to the schema.

How will it help you?

The schema has high discriminatory power and allows for the detection of markers specific for subtypes or outbreaks, thus enabling more powerful classification and outbreak definition tools. Together with BioNumerics and our powerful cloud based Calculation Engine, it completes a high-throughput environment that enables a faster and a lot more straightforward analysis of whole genome sequencing data

for *B. cereus*. The Calculation Engine's quality-controlled de novo assembly possibilities allow you to easily assemble whole genome sequencing data without the need of local computing power. Moreover, the two allele detection procedures (assembly-based and assembly free) allow you to perform fast and reliable allele calling for e.g. cluster detection which can be combined with whole genome SNP analysis to obtain the utmost resolution within your sample comparisons.

The whole-genome multi-locus sequence typing schema for *B. cereus* has been tested, validated and approved by our microbiologists on several public datasets.

With turnaround times of less than 30 minutes per sample and simultaneous processing of multiple samples, the power of high-performance computing is brought to your desktop with only a few clicks.

Simply request a Calculation Engine project to get started. We have wgMLST tutorials available online.

Don't have a BioNumerics license yet? Try it for FREE for one month.

INTERESTED?

Request a calculation engine project today to get started:

